arXiv:1211.3903v1 [math.OA] 16 Nov 2012

Birkhoff's ergodic theorem in operator algebras

Anilesh Mohari

The Institute of Mathematical Sciences,

Taramani, CIT Campus, Chennai-600113

E-mail:anilesh@imsc.res.in

Abstract

Let $\mathcal{M}$ be a $\sigma-$finite von-Neumann algebra with a separable pre-dual and α be an automorphism on $\mathcal{M}$ with a faithful normal state ϕ. It is well known that there exists a norm-one projection $E : \mathcal{M} \rightarrow \mathcal{M}$ so that $s_n(x) = \frac{1}{n}\sum_{0 \le k \le n-1} \alpha^k(x) \rightarrow E(x)$ as $n \rightarrow \infty$ for all $x \in \mathcal{M}$ in the weak* topology induced by it's pre-dual Banach space $\mathcal{M}_*$ where E is the norm one projection on von-Neumann sub-algebra $\mathcal{N} = \{x \in \mathcal{M} : \alpha(x) = x\}$. We prove in the present short paper that

$$\lim_{n \rightarrow \infty} ||\psi \circ s_n - \psi \circ E|| \rightarrow 0$$

as $n \rightarrow \infty$ where $\psi \in \mathcal{M}_*$ and $||.||$ is the Banach space norm of $\mathcal{M}_*$. We also extend the scope of the result to a normal unital Markov map $\tau : \mathcal{M} \rightarrow \mathcal{M}$ which admits a faithful normal invariant state ϕ. The result includes a proof for a second countable locally compact amenable group G acting on $\mathcal{M}$.

1 Introduction:

Let $\mathcal{H}$ be a complex Hilbert space and $\mathcal{B}(\mathcal{H})$ be the set of bounded operators on $\mathcal{H}$. $\mathcal{B}(\mathcal{H})$ is the Banach space dual of the trace class operators $\mathcal{T}(\mathcal{H})$. A $*$-sub-algebra $\mathcal{M}$ of $\mathcal{B}(\mathcal{H})$ is called von-Neumann algebra if $\mathcal{M}$ is closed in weak* topology on $\mathcal{B}(\mathcal{H})$. Given a von-Neumann algebra $\mathcal{M}$, the pre-dual Banach space $\mathcal{M}_*$ of $\mathcal{M}$ need not be separable in general and thus the norm closed unit ball in $\mathcal{M}$, which is compact in weak* topology, need not be metrizable and thus need not be sequentially compact i.e. a bounded sequence in $\mathcal{M}$ may not admit a convergent sub-sequence in the weak* topology. One usually avoid such a difficulties by considering $\mathcal{M}$ acting on a complex separable Hilbert space $\mathcal{H}$. For finer details on these questions we refer to Chapter 3 in [Tak].

Though any automorphism on $\mathcal{M}$ is normal and closed unit ball in the dual space $\mathcal{M}^*$ is compact in weak* topology, it may not ensure an invariant normal state for an automorphism on $\mathcal{M}$ and neither we can ensure even existence of an invariant state. This is not a surprising fact as these are well known when $\mathcal{M}$ is a commutative von-Neumann algebra such as $l_\infty(\mathbb{Z})$ and automorphism is the shift. There exists no invariant state on $l_\infty(\mathbb{Z})$ for a shift. Difficulty lies with the fact that $\mathcal{M}$ is not separable as a C^*-algebra in most interesting cases which includes example such as $\mathcal{B}(\mathcal{H})$, where $\mathcal{H}$ is a complex separable Hilbert space. However there are many well known situation, like in ergodic theory on measure space, in quantum theory as well, where an automorphism admits an invariant normal state. Most interesting example arises in the GNS space associated with a C^*-dynamical systems associated with invariant state. We will discuss briefly C^*-dynamical systems at the end of this paper.

We call a triplet $(\mathcal{M}, \alpha, \phi)$ a quantum dynamical system where $\mathcal{M}$ is a von-

Neumann algebra and α is an automorphism with a faithful normal invariant state ϕ. The assumption that ϕ is faithful does not put a strong restriction as otherwise we can restrict our attention to the corner von-Neumann algebra $p_\phi \mathcal{M} p_\phi$ where $p_\phi \in \mathcal{M}$ is the support projection of the normal state ϕ as p_ϕ is an invariant projection for α. In such a case the corner von-Neumann algebra is a $\sigma-$finite von-Neumann algebra. We will also assume without loss of generality that $\mathcal{M}$ be a $\sigma-$finite von-Neumann algebra acting on a Hilbert space $\mathcal{H}$ which is equivalent to existence of a faithful normal state on $\mathcal{M}$ [BR vol-1].

Let $(\mathcal{M}, \alpha, \phi)$ be a quantum dynamical system. It is quite some time now that it is known [Fr] that

$$s_n(x) = \frac{1}{n} \sum_{1 \le k \le n} \rightarrow E(x) \tag{1.1}$$

as $n \rightarrow \infty$ in weak* topology i.e. $\sigma-$weak operator topology i.e. for each $\psi \in \mathcal{M}_*$ $\psi(s_n(x)) \rightarrow \psi(E(x))$ for all $x \in \mathcal{M}$ as $n \rightarrow \infty$. Given a sequence of weakly convergent normal states ψ_n on $\mathcal{M}$, the sequence need not be strongly convergent i.e. need not be Cauchy in Banach space norm of $\mathcal{M}_*$. Such a statement however true if $\mathcal{M}$ is a type-I von-Neumann algebra with centre completely atomic. In other situation it is known to be false [De]. Thus the general theory of strong convergence no way ensures Birkhoff theorem which says that

$$||\psi \circ s_n - \psi \circ E|| \rightarrow 0 \tag{1.2}$$

as $n \rightarrow \infty$ for all element $\psi \in \mathcal{M}_*$. Interesting point here that faithful property of the invariants state allows us to avoid assuming sequential compactness of the closed unit ball of $\mathcal{M}$ in weak* topology. Thus the result that we prove here is a bit of surprise though Birkhoff point-wise ergodic theorem makes little sense in non-commutative framework of operator algebras.

The paper is organized as follows: In our next section we enlarge our domain to quantum dynamical semi-group of completely positive normal map on $\mathcal{M}$ after a short review on completely positive map on C^* algebra. In section 3 we will include Birkhoff theorem for a second countable locally compact amenable group action G on a von-Neumann algebra $\mathcal{M}$.

2 Quantum dynamical semi-group:

Let $\mathcal{A}$ be an unital C^* algebra. A map τ on $\mathcal{A}$ is called positive if $\tau(\mathcal{A}_+) \subseteq \mathcal{A}_+$, where $\mathcal{A}_+$ is the cone consists of the positive elements of $\mathcal{A}$ i.e. $\mathcal{A}_+ = \{x^*x : x \in \mathcal{A}\}$. τ is called n-positive if

$$\tau \otimes I_n : (x^i_j) \to (\tau(x^i_j))$$

is also positive on $M_n(\mathcal{A})$, $(n \times n)$ matrices with elements in $\mathcal{A}$. τ is called completely positive if $\tau \otimes I_n$ is positive for each $n \geq 1$. By Gelfand's theorem without loss of generality we may assume that $\mathcal{A} \subseteq \mathcal{B}(\mathcal{H})$, where $\mathcal{H}$ is a Hilbert space over field of complex numbers. It is well know [St] that such a completely positive map $\tau : \mathcal{A} \to \mathcal{B}(\mathcal{H})$ admits a canonical representation given by

$$\tau(x) = V^*\pi(x)V$$

where $\pi : \mathcal{A} \to \mathcal{B}(\mathcal{K})$ is representation of $\mathcal{A}$ in $\mathcal{B}(\mathcal{K})$, $\mathcal{K}$ is a Hilbert space over field of complex numbers and V is a bounded operator from $\mathcal{H} \to \mathcal{K}$. Further τ is unital if and only if V is an isometry. In this paper we will consider only unital maps.

Given a unital $2-$positive map τ on $\mathcal{A}$ we have Kadison-Schwarz inequality

$$\tau(x^*)\tau(x) \leq \tau(x^*x) \tag{2.1}$$

and further if equality hold for x then

$$\tau(x^*)\tau(y) = \tau(x^*y) \tag{2.2}$$

for all $y \in \mathcal{A}$. This shows that $\{x : \tau(x^*)\tau(x) = \tau(x^*x),\ \tau(x)\tau(x^*) = \tau(xx^*)\}$ is C^*-sub algebra of $\mathcal{A}$.

We take $I\!\!T$ to be either $Z\!\!\!Z$, the set of integers or $I\!\!R$ the set of real numbers. $I\!\!T_+$ be set of positive numbers of $I\!\!T$. Let $\tau_t : t \geq 0$ be a semi-group of completely positive maps on von-Neumann algebra $\mathcal{M}$ with $\tau_0 = I$. We assume that for each $t \geq 0$, τ_t is a normal map i.e. $l.u.b.\tau_t(x_\alpha) = \tau_t(l.u.b.x_\alpha)$ for any increasing net $x_\alpha \in \mathcal{M}_+$ bounded from above where $l.u.b.$ is the least upper bound and $t \rightarrow \tau_t(x)$ is continuous in $\sigma-$weak operator topology when $I\!\!T = I\!\!R$. Let ϕ be a faithful normal τ-invariant state on $\mathcal{M}$ i.e.

$$\phi(\tau_t(x)) = \phi(x)$$

for all $x \in \mathcal{M}$ and $t \geq 0$. The result that follows in this section has a ready generalization to a faithful weight [Ta] which we avoid as it makes very little qualitative improvement of our results once we incorporate Radon-Nykodym theorem for weights. One interesting observation here that once we know a positive map τ admits a faithful normal invariant state, then τ is automatically normal.

Lemma 2.1: Let ϕ be a faithful normal invariant state for a completely positive unital map $\tau : \mathcal{M} \rightarrow \mathcal{M}$. We identify $\mathcal{M}$ with $\pi_\phi(\mathcal{M})$ where $(\mathcal{H}_\phi, \pi_\phi, \zeta_\phi)$ is the GNS space associated with ϕ, $\phi(x) =< \zeta_\phi, \pi_\phi(x)\zeta_\phi >$ and ζ_ϕ is a cyclic and separating unit vector for $\pi_\phi(\mathcal{M})$ in $\mathcal{H}_\phi$. Then there exists a completely positive map $\tau' : \pi_\phi(\mathcal{M})' \rightarrow \pi_\phi(\mathcal{M})'$ so that

$$< \tau'(y')\zeta_\phi, x\zeta_\phi >=< y'\zeta_\phi, \tau(x)\zeta_\phi > \tag{2.3}$$

for all $x \in \pi_\phi(\mathcal{M})$ and $y' \in \pi_\phi(\mathcal{M})'$. Further

$$|\psi_{y'_1,y'_2}(s_n(x) - s_m(x))| \le ||(s'_n - s'_m)(y^*_1 y_2)\zeta_\phi||||x\zeta_\phi|| \le ||(s'_n - s'_m)((y')^*_1 y'_2)\zeta_\phi||\ ||x|| \tag{2.4}$$

where $\psi_{y'_1,y'_2}(x) =< y'_1\zeta_\phi, xy'_2\zeta_\phi >$ and s_n, s'_n denotes n^{th} mean associated with the maps τ and τ' respectively.

Proof: First part of the statement is not new and details have been worked out in section 8 in [OP]. For the sack of completeness we recall essential points omitting the details. Existence part is a standard result which follows as a simple application of Dixmier's fundamental lemma [Di] for normal state ψ satisfying $\psi \le \lambda\phi$ on $\mathcal{M}_+$ for some $\lambda > 0$ admits a representation $\psi(x) =< y'\zeta_\phi, x\zeta_\phi >$ where y' is non-negative element in the commutant $\pi_\phi(\mathcal{M})'$ of $\pi_\phi(\mathcal{M})$ and such an element is unique. For any non-negative element $y' \in \pi_\phi(\mathcal{M})'$ we consider non-negative normal functional $\psi_{y'}(x) =< y'\zeta_\phi, \tau(x)\zeta_\phi >$ and check that $\psi_{y'} \le ||y'||\phi$ and thus by Dixmier's lemma we get an unique non-negative element $\tau'(y') \in \pi_\phi(\mathcal{M})'$ so that (2.3) is satisfied. We can extend now the definition of $\tau'(y')$ for an arbitrary self adjoint element by linearity using the decomposition $y' = y'_+ - y'_-$ with $y'_+ y'_- = 0$. For an arbitrary element y we use $\pi_\phi(\mathcal{M})' = \pi_\phi(\mathcal{M})'_h + i\pi_\phi(\mathcal{M})'_h$, $\pi_\phi(\mathcal{M})'$ being a $*$-algebra. It is simple to check that (2.3) hold for any $y' \in \pi_\phi(\mathcal{M})', x \in \pi_\phi(\mathcal{M})$.

That τ' takes non-negative element of $\mathcal{M}'$ to non-negative elements follows by our construction and also note that $\tau'(I) = I$ since $\phi(\tau(x)) = \phi(x)$ and ϕ is an invariant state for τ'. The vector state $y' \rightarrow < \zeta_\phi, y'\zeta_\phi >$ being normal faithful on $\mathcal{M}'$, invariant property for τ' will ensure normal property of τ' from normality of the vector state.

That n-positivity of τ' will follow from $n-$positivity of τ once we use modular relation to rewrite (2.3) as follows:

$$\phi(\tilde{\tau}(y)\sigma_{\frac{i}{2}}(x)) = \phi(\sigma_{-\frac{i}{2}}(y)\tau(x)) \tag{2.5}$$

for all $x, y \in \mathcal{M}$ those are analytic with respect to the modular group $\sigma = (\sigma_t)$ where $\sigma_t(x) = \Delta_\phi^{it} x \Delta_\phi^{-it} : t \in I\!R$ on $\mathcal{M}$ associated with ϕ where

$$\tilde{\tau}(y) = \mathcal{J}_\phi \tau'(\mathcal{J}_\phi y \mathcal{J}_\phi)\mathcal{J}_\phi$$

and $\Delta_\phi, \mathcal{J}_\phi$ are modular operator associates with Tomita's conjugate action

$$S_0 : x\zeta_\phi \to x^*\zeta_\phi, x \in \mathcal{M}.$$

We use modular relation

$$\phi(x^*y) = \phi(y^*(\phi)x(\phi)) \tag{2.6}$$

for two analytic elements x, y where we write $x(\phi) = \sigma_{\frac{i}{2}}(x)$ for any analytic element x for the modular group.

Thus we check for analytic elements $1 \leq x_i, y_i \leq n$ we have that $\phi(y_i^*\tilde{\tau}(x_i^*x_j)y_j) = \phi(x_j(\phi)^*\tau(y_j(\phi)^*y_i(\phi))x_i(\phi))$ and thus $\sum_{i,j}\phi(y_i^*\tilde{\tau}(x_i^*x_j)y_j) = \sum_{i,j}\phi(x_j(\phi)^*\tau(y_j(\phi)^*y_i(\phi))x_i(\phi)) \geq 0$ by n-positivity of τ. By faithful property we get $y_i^*\tilde{\tau}(x_i^*x_j)y_j) \geq 0$. Since analytic elements are dense in weak* topology we also have $[\mathcal{M}_a\zeta_\phi] = [\mathcal{M}\zeta_\phi] = \mathcal{H}_\phi$. Hence $((\tilde{\tau}(x_i^*x_j)))$ is a non-negative definite element in $\mathcal{M} \otimes M_n(I\!\!C)$. Thus $\tilde{\tau} \otimes I_n$ takes non-negative elements of the form $((x_i^*x_j)) : x_i \in \mathcal{M}_a)$ into non-negative elements of $M_n(\mathcal{M})$. Such elements being dense in $M_n(\mathcal{M})_+$, we conclude the result by normality of $\tilde{\tau} \otimes I_n$.

We can avoid use of Tomita's theorem while proving n-positivity of τ' as follows. That τ' is positive follows from Dixmier's lemma. For n positivity of τ', we consider the faithful normal state $\phi_n = \phi \otimes tr_n$ on $M_n(\pi_\phi(\mathcal{M}))$, $n \times n$ matrices with entries in $\pi_\phi(\mathcal{M})$ and identify with it standard form i.e. GNS space, where tr_n is the normalize trace on $M_n(\mathbb{C})$. Now we consider the map $\tau_n = \tau \otimes I_n$ in the GNS space associated faithful normal invariant state ϕ_n and check that the dual map of Dixmier is given by $\tau'_n = \tau' \otimes I_n$ in the commutant of $M_n(\pi_\phi(\mathcal{M}))' = M_n(\pi_\phi(\mathcal{M})')$. Thus positivity property of τ'_n follows from that of τ_n.

The inequality (2.4) is fairly obvious by Cauchy-Schwarz inequality. ■

Lemma 2.2: Let $(\mathcal{M}, \tau, \phi)$ be as in Lemma 2.1. The map $T : x\zeta_\phi \to \tau(x)\zeta_\phi$ has a unique contractive extension on $\mathcal{H}_\phi$ and

$$s_n(T) = \frac{1}{n} \sum_{1 \le k \le n-1} T^k \to P \tag{2.7}$$

in strong operator topology where P is the projection on the closed subspace $\{f : Tf = f\}$ which is equal to the closed space generated by $\{x\zeta_\phi : x \in \mathcal{N}\}$ where $\mathcal{N} = \{x \in \mathcal{M} : \tau(x) = x\}$. Further

$$s_n(x) \to E(x) \tag{2.8}$$

in weak* topology of $\mathcal{M}$.

Proof: Once again first part is a well known consequence of Kadison-Schwarz inequality for 2-positivity map τ which says that

$$\tau(x^*)\tau(x) \le \tau(x^*x) \tag{2.9}$$

for all $x \in \mathcal{M}$ and further equality hold in (2.9) for x if and only if

$$\tau(x^*y) = \tau(x^*)\tau(y) \tag{2.10}$$

for all $y \in \mathcal{M}$. Thus we get $||Tx\zeta_\phi|| \le ||x\zeta_\phi||$ by invariance property. Let P be the orthogonal projection on the closed subspace $\{f : Tf = f\}$. Then $(1-P)TP = 0$. For any $f \in \mathcal{H}_\phi$, with $Tf = f$, we also have $||T^*f - f||^2 =$

$||T^*f||^2 - < T^*f, f > - < f, T^*f > + ||f||^2 \leq 0$ as T^* is a contraction and $Tf = f$. Thus we get $T^*f = f$. Thus we also have $(1-P)T^*P = 0$ and so $TP = PT$. It is simple to check also that $s_n(T)(I-T) = \frac{1}{n}(I - T^n)$ and so $||s_n(T)(I-T)|| \leq \frac{2}{n} \rightarrow 0$ as $n \rightarrow \infty$. In particular we get $s_n(T)f \rightarrow 0$ in strong operator topology as $n \rightarrow \infty$ for any $f \in R(I-T) = N(I-T^*)^{\perp} = I - P$ i.e. $s_n(T) \rightarrow P$ in strong operator topology as $n \rightarrow \infty$. The line of argument that we use here can be traced back to P. R. Halmos [Ha].

We apply compactness property for the bounded net $s_n(x)$ as $||s_n(x)|| \leq ||x||$ to claim that it has unique limit point. Let $x^1_{\infty}, x^2_{\infty}$ be two such limit points. If so then $x^1_{\infty}\zeta_{\phi} = x^2_{\infty}\zeta_{\phi}$ as $s_n(x)\zeta_{\phi} \rightarrow Px\zeta_{\phi}$ in strong operator topology. Now by separating property of ζ_{ϕ} for $\mathcal{M}$, we conclude that $x^1_{\infty} = x^2_{\infty}$. Limit point being unique we conclude that $s_n(x) \rightarrow x_{\infty}$ for some x_{∞} in weak* topology. It is now a routine work to check that the map $x \rightarrow E(x) = x_{\infty}$ is completely positive unital map preserving ϕ. Thus E is also normal. As $Px_{\infty}\zeta_{\phi} = x_{\infty}\zeta_{\phi}$ i.e. $\tau(x_{\infty}) = x_{\infty}$ by separating property and so $E(x) \in \mathcal{N}$. That $E^2 = E$ is obvious and so is the bi-module property of E [Um] i.e. $yE(x)z = E(yxz)$ for all $y, z \in \mathcal{N}$ and $x \in \mathcal{M}$ which follows from (2.10) applied to τ_n

$$y\tau_n(x)z = \tau_n(yxz)$$

for all $y, z \in \mathcal{N}$ and $x \in \mathcal{M}$. ■

Theorem 2.3: Let $(\mathcal{M}, \tau, \phi)$ be a completely positive unital map with a faithful normal invariant state on $\mathcal{M}$. Then there exists a norm one projection $E : \mathcal{M} \rightarrow \mathcal{N}$ onto $\mathcal{N}$ so that

$$||\psi \circ s_n - \psi \circ E|| \rightarrow 0 \tag{2.11}$$

as $n \rightarrow \infty$ for any $\psi \in \mathcal{M}_*$.

Proof: Lemma 2.2 is saying that (2.11) is true for all normal state $\psi = \psi_{y_1,y_2}$. Since ζ_ϕ is also cyclic for $\pi_\phi(\mathcal{M})'$, the class of normal functional $\{\psi_{y_1,y_2} : y_1, y_2 \in \pi_\phi(\mathcal{M})'\}$ is total in the Banach space $\mathcal{M}_*$ in strong topology, we conclude same holds for any element $\mathcal{M}_*$. ■

Theorem 2.4: Let $(\mathcal{M}, \tau_t, \phi)$ be a semi-group of completely positive unital map with a faithful normal invariant state on $\mathcal{M}$. Then there exists a norm one projection $E : \mathcal{M} \to \mathcal{N}$ onto $\mathcal{N}$ so that

$$||\psi \circ s_\lambda - \psi \circ E|| \to 0 \tag{2.12}$$

as $\lambda \to 0$ for any $\psi \in \mathcal{M}_*$ where

$$s_\lambda = \lambda \int e^{-\lambda t} \tau_t dt$$

and E is the norm one normal projection onto $\mathcal{N} = \{x : \tau_t(x) = x,\ t \geq 0\}$.

Proof: Proof goes along the same line that of discrete time dynamics [Fr]. We indicate here a proof leaving the details to reader. We set contractive operator

$$s_\lambda(T) = \lambda \int e^{-\lambda t} T_t dt$$

on $\mathcal{H}_\phi$ where $T_t x \zeta_\phi = \tau_t(x)\zeta_\phi$ is the strongly continuous semi-group of contraction on $\mathcal{H}_\phi$. As $\lambda \to 0$, $s_\lambda \to P$ in strong operator topology follows along the same line where P is the projection on the closed subspace $\{f : T_t f = f : t \geq 0\}$. Thus we can use strong convergence $s'_\lambda(T') \to P'$ as $\lambda \to 0$, where P' is the projection on the closed subspace $\{f : T'_t f = f : t \geq 0\}$ where $T'_t y' \Omega = \tau'(y')\Omega,\ y' \in \mathcal{M}'$. Once again same argument shows that $P' = P$ to conclude that $||\psi_{y'_1,y'_2}(s_\lambda - s_\mu)|| \to 0$ in $\mathcal{M}_*$ as $\lambda, \mu \to 0$. Once more using sequential weak* compactness of $\mathcal{M}$ and duality argument, we conclude that $||\psi \circ s_\lambda - \psi \circ E||$ as $\lambda \to 0$ for all $\psi \in \mathcal{M}_*$. ■

Note that for E to be a normal norm one projection from $\mathcal{M}$ on $\mathcal{N}$ in above theorems, we only require τ to be $2-$positivity to ensure Kadison-Schwarz inequality (2.9). For a normal invariant state ϕ for τ we have $\tau_t(p_\phi) \ge p_\phi$ for all $t \ge 0$ and $\tau^\phi_t(x) = p_\phi \tau_t(p_\phi x p_\phi) p_\phi$ is a one parameter semi-group of completely positive ($2-$positive if τ is so) contractive unital map on the corner von-Neumann algebra $\mathcal{M}_\phi = p_\phi \mathcal{M} p_\phi$. In such a case if $\mathcal{N} = \{x : \tau_t(x) = x\}$ we have $\tau_t(p_\phi) \uparrow I$ as $t \to \infty$ and $s_\lambda(x) \to \phi_0(x)$ in weak* topology as $\lambda \to 0$ [Mo2]. Thus going along the same line we have $||\psi \circ s_\lambda - \phi|| \to 0$ as $n \to \infty$ if $\mathcal{N} = \{x : \tau_t(x) = x : t \ge 0\} = \{zI : z \in \mathbb{C}\}$ for any normal state ψ on $\mathcal{M}$.

In the next section we focus our interest to a locally compact second countable amenable group G acting on a von-Neumann algebra $\mathcal{M}$ with a separable pre-dual space and prove Birkhoff theorem.

3 Birkhoff's ergodic theorem for amenable group:

A locally compact group G is called amenable of for any compact $K \subset G$ and $\delta > 0$ there exists a compact subset $F \subset G$ so that $|F \Delta KF| < \delta |F|$ where we use $|.|$ to denote the left Haar measure on G. Such a set F is called $(K,\delta)-$invariant. An infinite sequence $F_1, F_2, ...$ of compact subsets of G will be called *Følner sequence* if for every compact K and $\delta > 0$, F_n is (K,δ)-invariant for all $n \ge N(K,\delta)$, where $N(K,\delta)$ is an integer depending on K and δ.

Let G be a locally compact second countable amenable group acting from left bi-measurably on a probability space $(X, \mathcal{F}, \mu)$ such that

$$\mu \circ g^{-1}(E) = \mu(E)$$

for all $g \in G$ and $E \in \mathcal{F}$. For an element $f \in L^1(x, \mathcal{B}, \mu)$, we set $s(F, f)$ to denote the average

$$s(F, f)(x) = \frac{1}{|F|} \int_F f(gx) dm(g)$$

where dm is a left-invariant Haar-measure on G.

A basic problem in ergodic theory is to make a wise choice for a Følner sequence so that

$$s(F_n, f) \to \bar{f}(x) \text{ a.e.}$$

and also in norm topology of $L^1(X, \mathcal{B}, \mu)$ as $n \to \infty$ where $\bar{f}(x) = E(f|\mathcal{F}_e)(x)$ is the conditional expectation f on the σ-field of G-invariant sets i.e.

$$\mathcal{F}_e = \{E : \mu(E \Delta g^{-1}(E)) = 0, \ \forall g \in G\}$$

A. Shulman [Sh] introduced in his thesis such an useful concept to prove an L^2-version of ergodic theorem [Sh] which played a key role in Lindenstrauss L^1 version of ergodic theorem [Li]. We briefly recall [Sh] that a Følner sequence $\{F_n : n \geq 1\}$ of compact subsets of G is called tempered if there exists a constant $C > 0$ such that

$$|\bigcup_{k<n} F_k^{-1} F_n| < C|F_n|$$

for all $n \geq 1$. Lindenstrauss [Li] also proven that there exists a sub-sequence of a Følner sequence which satisfies A. Shulman's tempered condition.

A. Shulman [Sh] proved that for each $f \in L^2(X, \mathcal{B}, \mu)$

$$s(F_n, f) \to \bar{f}$$

in L^2 whenever F_n is a tempered Følner sequence. In the following we put his result in an abstract set up.

Theorem 3.1: [Sh] Let G be a second countable locally compact amenable group and F_n be a tempered Følner sequence for G. Let $g \to u_g$ be a unitary representation on a Hilbert space $\mathcal{H}$ so that the map $g \to < f, u_g h >$ be continuous for any two elements $f, h \in \mathcal{H}$. Then

$$s(F_n, f) \to Pf \tag{3.1}$$

in strong operator topology as $n \to \infty$ where P is the projection on the closed subspace $\{f : u_g f = f, \ \forall g \in G\}$ and

$$s(F, f) = \frac{1}{|F|} \int_F u_g f dm(g)$$

Proof: It goes along the same line [Sh] once we notice the proof does not use explicit form of the group action on the Hilbert space $L^2(X, \mathcal{B}, \mu)$. We skip the details. ■

Theorem 3.2: Let G, F_n be as in Theorem 3.1. Let $g \to \alpha_g$ be a group of $*$-automorphism on $\mathcal{M}$ such that $g \to \psi(\alpha_g(x))$ is a measurable function for each $x \in \mathcal{M}$ and $\psi \in \mathcal{M}_*$ and ϕ be a faithful normal invariant state for α_g. Then for any $\psi \in \mathcal{A}_*$

$$||\psi \circ s_n - \psi \circ E|| \to 0$$

as $n \to \infty$ where E is the normal norm one projection from $\mathcal{M}$ to

$$\mathcal{N} = \{x \in \mathcal{M} : \alpha_g(x) = x\}$$

Proof: We set unitary representation $g \to u_g : x\zeta_\phi = \alpha_g(x)\zeta_\phi, \ x \in \mathcal{M}$ and $s_n(u) = \frac{1}{|F_n|} \int_{F_n} u_g dm(g)$. By Theorem 3.1 we have $s_n \to P$ in strong operator topology where P is the projection on invariant vectors $\{f : u_g f = f : \ g \in G\}$. We claim that P is the close span of the vectors in $\{x\zeta_\phi : x \in \mathcal{N}\}$ where $\mathcal{N} = \{x : \alpha_g(x) = x; g \in G\}$. To that end we consider

$$s_n(x) = \frac{1}{|F_n|} \int_{F_n} \alpha_g(x) dm(g)$$

and we once again we conclude that $s_n(x) \to E(x)$ in weak* topology by weak* compactness of the unit ball to some element $E(x)$ and faithfulness of the normal state. So in particular we have $E(x)\zeta_\phi \in P$. Thus we have $E(x)\zeta_\phi = \alpha_g(E(x))\zeta_\phi$ for all $g \in G$. Thus $E(x) \in \mathcal{N}$ by separating property of ζ_ϕ. Still we have not shown that P is the linear span of $\{u_g f = f : f \in \mathcal{H}_\phi\}$. One way inclusion is trivial. For the reverse direction, let f be an element such that $u_g f = f$ for all $g \in G$ and orthogonal to $\{x\zeta_\phi : x \in \mathcal{N}\}$. Then $< f, E(x)\zeta_\phi >= 0$ for all $x \in \mathcal{M}$. However we also have $< f, E(x)\zeta_\phi >= \lim_{n\to\infty} < f, s_n(x)\zeta_\phi >=< f, x\zeta_\phi >$. ζ_ϕ being cyclic we conclude that $f = 0$. This shows the equality of those two subspaces.

Now we use (2.4) with $x \to s_n(x)$, which is a completely positive map on $\mathcal{M}$, to conclude

$$||\psi_{y_1',y_2'} \circ s_n - \psi_{y_1',y_2'} \circ s_m|| \le ||(s_n' - s_m')((y_2')^* y_1)\zeta_\phi|| \to 0$$

as $m, n \to \infty$ as $s_n'(u) \to P$ in strong operator topology by Theorem 3.1 where $s_n'(u) = \frac{1}{|F_n|} \int u_{g^{-1}} dm(g)$ on $\mathcal{H}_\phi$. Thus the result follows since the set $\{\psi_{y_1',y_2'} : y_1', y_2' \in \mathcal{M}'\}$ is total in $\mathcal{M}_*$ in the Banach space topology. ■

Before we end this paper we briefly review classical situation. Let G be as in Theorem 3.2. G be acting on a probability space $(X, \mathcal{B}, \mu)$ bi-measurably i.e. $(g, x) \to gx$ is a measurable map so that for each g the action $x \to gx$ is measure preserving. Let $\alpha_g : L^\infty(X, \mathcal{F}, \mu) \to L^\infty(X, \mathcal{F}, \mu)$ be the automorphism defined by

$$\alpha_g(h)(x) = h(gx), \ x \in X$$

Null sets being preserved by each g, the map is well defined. $L^\infty(X,\mathcal{B},\mu)$ is identified with $\{M_h : h \in L^\infty(X,\mathcal{B},\mu)$ where M_h is the multiplication function on $L^2(X,\mathcal{B},\mu)$ defined by

$$M_h(f)(x) = h(x)f(x),\ x \in X$$

a element $\psi \in \mathcal{M}_*$ is written as $\psi(x) = \sum_{k\ge 1} < f_n, xg_n >$ where $f_n, g_n \in L^2(X,\mathcal{B},\mu)$ with $\sum_n ||f_n||^2 < \infty$ and $\sum_{n\ge 1} ||g_n||^2 < \infty$. Thus we may identify ψ with $\psi(x) = \sum_n f_n(x)\overline{g_n(x)} \in L^1(X,\mathcal{B},\mu)$. Note at this point that $\psi(M_h) =< \psi, h >= \int_X \overline{\psi(x)}h(x)d\mu(x)$. In our analysis we have assumed that closed unit ball $\mathcal{M}$ is sequentially compact. Here $\mathcal{M} = L^\infty(X,\mathcal{B},\mu)$ is automatically $\sigma-$finite von-Neumann algebra as μ is a faithful state on $\mathcal{M}$. However sequential compactness of closed unit ball of $\mathcal{M}$ in weak* topology is not so obvious unless we assume $\mathcal{M}_*$ is separable. Proposition 2.1 with commutative $\mathcal{M}$ appeared in the literature [Ne] while studying Hunt's notion of time reverse Markov process. The net gain that we have got in the classical set up that we could arrive at Birkhoff's L^1- convergence theorem without proving ‘ maximal ergodic theorem ’ [Wa,Li] and Birkhoff L^1 convergence is intimately connected to von Neumann's L^2 ergodic theorem for dual map. It is not clear however how one can arrive at Birkhoff's maximal ergodic theorem by a functional analytic method.

We refer to monographs [BR] or [Pe] for interesting examples that appear in C^*-dynamical systems. In the following we briefly indicate standard set up for a C^*-dynamical system. Let G be a second countable locally compact amenable group and $x \to \alpha_g(x),\ g \in G$ be a group of $*$-automorphism on a unital C^*-algebra $\mathcal{A}$. We assume for each $x \in \mathcal{A}$ the map $g \to \alpha_g(x)$ is continuous. We may assume sequential weak* compactness of the norm closed unit ball of dual space $\mathcal{A}^*$ of $\mathcal{A}$ for existence of an invariant state for α_g. Let ϕ be a G-invariant state for $\mathcal{A}$. It is simple to check that the map $\pi_\phi(x) \to \pi_\phi(\alpha_g(x))$ extends to an automorphism on $\mathcal{M} = \pi_\phi(\mathcal{A})''$ and for each $x \in \mathcal{M}$, the map $g \to \pi_\phi(\alpha_g(x))$ is continuous in $\sigma-$ weak operator topology where $(\mathcal{H}_\phi, \pi_\phi, \zeta_\phi)$ is the GNS space of $(\mathcal{A}, \phi)$. A state ϕ on $\mathcal{A}$ is called faithful if $\phi(x) = 0$ for $x \geq 0$ ensures that $x = 0$. In such a case the vector state by ζ_ϕ is a faithful normal invariant state for the induced group of automorphisms (α_g) on $\mathcal{M}$. The extremal elements in the convex set of β-KMS state for a translation invariant Hamiltonian on one dimensional quantum spin $\mathcal{A} = \otimes_{\mathbb{Z}} M_d(\mathbb{C})$ are known examples that are faithful. Such a state is also known to be a factor state and ergodic for the translation action. For more explicit examples we refer to [BR,OP,NeS,BE]. Duality relation (2.3) has been explored in greater details in a series of papers [Mo1-3].

REFERENCES

- [BR] Bratteli, O., Robinson, D.W. : Operator algebras and quantum statistical mechanics, I,II, Springer 1981.
- [BE] Bratteli, O., Evans, David E. . Dynamical semigroups commuting with compact abelian actions. Ergodic Theory and Dynamical Systems, 3 (1983), pp 187-217.
- [Di] Dixmier, J.: von Neumann algebras, North Holland 1981.
- [De] Dell'Antonio, G. F.: On the limits of sequences of normal states. Comm. Pure Appl. Math. 20 (1967), pp 413-429.
- [Fr] Frigerio, A.: Stationary states of quantum dynamical semigroups. Comm. Math. Phys. 63 (1978), no. 3, 269-276.
- [Lin] Lindenstrauss, E.: Point-wise theorems for amenable groups. Invent. Math. 146 (2001), no. 2, pp 259-295.

- [Ha] Halmos, P.R.: Lecture on ergodic theory, Chelsea Pub. Co. New York 1956.
- [Mo1] Mohari, A.: Markov shift in non-commutative probability, Jour. Func. Anal. 199 (2003) 189-209.
- [Mo2] Mohari, A.: Pure inductive limit state and Kolmogorov's property, J. Funct. Anal. vol 253, no-2, 584-604 (2007) Elsevier Sciences.
- [Mo3] Mohari, A.: Pure inductive limit state and Kolmogorov's property -II, http://arxiv.org/abs/1101.5961.
- [Ne] Nelson, E. The adjoint Markoff process. Duke Math. J. 25 (1958) 671-690.
- [NeS] Neshveyev, S.; Størmer, E. : Dynamical entropy in operator algebras. Springer-Verlag, Berlin, 2006.
- [OP] Ohya, M., Petz, D.: Quantum entropy and its use, Text and monograph in physics, Springer 1995.
- [Pe] Pedersen, G. K.: C*-algebras and their automorphism groups, London Mathematical Society. Monographs No. 14, Academic Press, 1979.
- [Sak] Sakai, S.: C^*-algebras and W^*-algebras, Springer 1971.
- [Sh] Schulman, A.: Maximal ergodic theorem in groups, Dep. Lit. NI-INTI, No.2184 (1988).
- [St] Stinespring, W. F.: Positive functions on C^*-algebras. Proc. Amer. Math. Soc. 6, (1955) 211-216.
- [Tak] Takesaki, M.: Theory of operator algebra -I, Springer 2001.

- [Um] Umegaki, H.: Conditional expectation in an operator algebra. Thoku Math. J. (2) 6, (1954) 177-181.
- [Wa] Walter, P.: An introduction to ergodic theory, Springer 1975.